

SIAST Palliser Campus

Mathematics

MAT 201

Lecture Notes and Examples

Unit 2

The Definite Integral

by Alan Dill and Robert G. Petry

published by the Department of Mathematics, SIAST Palliser Campus

Copyright © 2011 Alan Dill, Robert G. Petry.

Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.3 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled “GNU Free Documentation License”.

Permission is granted to retain (if desired) the original title of this document on modified copies.




History

- Original document produced in 2011 entitled “The Definitie Integral” written by Alan Dill and Robert G. Petry. Published by the Department of Mathematics, SIAST Palliser Campus. A transparent copy of this document is available via
<http://www.campioncollege.ca/about-us/faculty-listing/dr-robert-petry>

1 Area Under a Curve

There are a number of methods we can use for approximating the area under a curve between two given points. These include:


- sketching a triangle or rectangle of approximately the same area as the area under the curve and calculating the area
- subdividing the area under the curve into squares, counting the squares and multiplying by the area of one square (this is most easily done using squared paper)
- subdividing the area under the curve into rectangular panels of known width, whose height is equal to the height of the curve at some point between the sides of the rectangle, and summing the areas of the rectangles



The first of these methods is the least accurate, but is good for getting a quick estimate. The accuracy of the second and third methods increases as the number of squares or rectangular panels increases.

1.1 The Midpoint Method

When using the third of the above methods, we need to decide at which point in the panels to take the height of the curve as the height of the panel. It turns out that, in most cases, we will get the best approximation of the area under the curve by defining the height of each panel as the height of the curve (i.e. the value of the function) at the *midpoint* of the panel. By doing this we will, to a large extent, balance out the extra area of the parts of the panels extending above the curve with the missing area of the parts of the panels that fall below the curve.



We call this approach the *midpoint method*. It should be fairly obvious that, as we decrease the width and increase the number of rectangular panels, the accuracy of the estimate also increases.

In symbols, we can write the sum of the areas found using the midpoint method as follows:

$$A_{ab} \approx \sum_{i=1}^n f(x_i^*) \Delta x = f(x_1^*) \Delta x + \cdots + f(x_n^*) \Delta x \quad (1)$$

where $f(x_i^*)$ is the value of the function at the midpoint x_i^* of each panel, and Δx is the width of each panel, $\Delta x = \frac{b-a}{n}$. A_{ab} is the actual area under the curve $f(x)$ between $x = a$ and $x = b$.

Examples

1. Estimate each of the following areas by sketching a triangle or rectangle:
 - (a) the area under the curve $y = x^2$ from $x = 1$ to $x = 4$
 - (b) the area given in Problem 6, p. 879 of the text
2. Estimate the area under the curve $y = \sqrt{2x+1}$ from $x = 1$ to $x = 7$ by the midpoint method.
 - (a) using panels of width 2
 - (b) using panels of width 1

Reading:

Sec. 30-5, pp. 875-879

Problems:

Ex. 5, p. 879 # 1-9 (odd), 17, 18, 19, 20

2 The Definite Integral

If we are asked to approximate the area under the curve of a given mathematical function between two given x -values, we can inscribe rectangles between the curve and the x -axis and add the areas of these rectangles. If only a few such rectangles were used, we would only get a rough approximation of the area. If, however, we were to use a larger number of rectangles, the amount of “empty space” above the rectangles would be decreased, and we would get a better approximation of the area. (See the two diagrams in Figure 1, which illustrate this idea.)




Figure 1: Approximating the area under a curve.


If you imagine that the number of rectangles is increased without bound, the sum of the areas of the rectangles will approach a limiting value that is equal to the exact area under the curve.

In general, the area under a curve $y = f(x)$ is given by

$$A = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x$$

where

$$\Delta x = \frac{b-a}{n}$$




Let ΔA represent the area $PQSU$ under the curve.

$$\begin{aligned} A_{PQRU} &\leq \Delta A \leq A_{PQST} \\ y\Delta x &\leq \Delta A \leq (y + \Delta y)\Delta x \end{aligned}$$

Dividing by Δx gives

$$y \leq \frac{\Delta A}{\Delta x} \leq y + \Delta y$$



Taking the limit as Δx approaches zero, we get

$$y \leq \lim_{\Delta x \rightarrow 0} \frac{\Delta A}{\Delta x} \leq y$$

since, as $\Delta x \rightarrow 0$, $\Delta y \rightarrow 0$.

The middle term, which is “sandwiched” between two terms that both become equal to y in the limit, is, in fact, the derivative of the area with respect to x , i.e. dA/dx . Thus, we can write

$$\frac{dA}{dx} = y$$

In differential form this becomes what one would intuitively expect for the infinitesimal area dA of the rectangle of height y and infinitesimal width dx , namely

$$dA = y \, dx$$

Taking the integral of both sides, we get

$$\int dA = \int y \, dx$$

But the integral of dA is, by definition, A . Using the notation A_{ax} to represent the area between $x = a$ and any given value of x , we can write

$$A_{ax} = F(x) + C$$

where $F(x)$ is the antiderivative of the function y , and C is the constant of integration.

The area between $x = a$ and $x = a$ must, of course be zero. We can use this information to determine the constant C .

$$A_{aa} = F(a) + C = 0$$

Solving for C , we get

$$C = -F(a)$$

We can then write an expression for the area between $x = a$ and $x = b$:

$$A_{ab} = F(b) + C$$

Substituting $C = -F(a)$ we get

$$A_{ab} = F(b) - F(a)$$

This leads us to the following formal definition of the definite integral.

We define the *definite integral* of a function $y = f(x)$ as follows:

$\int_a^b f(x) \, dx = F(b) - F(a) \tag{2}$

where a and b are called the *limits of integration*. $F(b)$ and $F(a)$ are the values of the integral (or antiderivative) for $x = b$ and $x = a$ respectively.

As we have shown above, the definite integral is numerically equal to the area under the curve $y = f(x)$ between $x = a$ and $x = b$, i.e.

$$A_{ab} = \int_a^b f(x) \, dx = F(b) - F(a) \quad (3)$$

It should be noted that the area A_{ab} will be *negative* if the function lies below the x -axis (i.e. is negative) over the interval $[a, b]$.

Examples

1. Find the exact area under the curve $y = \sqrt{2x+1}$ from $x = 1$ to $x = 7$. (This is the same area that we estimated earlier by the midpoint method.)

Evaluate the following definite integrals. Give the answers in exact form, and in approximate form to 3 significant digits.

2. $\int_2^3 \frac{1}{x^2} \, dx$

3. $\int_{-1}^2 (x^2 - 7)x \, dx$

4. $\int_0^1 \frac{x \, dx}{\sqrt{2-x^2}}$

5. $\int_{-1.6}^{0.7} \sqrt[3]{1-x} \, dx$

6. $\int_1^e \frac{dx}{x}$

7. $\int_0^2 \frac{dx}{8-3x}$

Additional Examples (Text, p. 875)

- Problem 6
- Problem 8
- Problem 12

Reading:

Secs. 30-4, 30-6; pp. 873-875, 880-882

Problems:

Ex. 4, p. 875 # 1-14 (all); Ex. 6, p. 883 # 1-12 (all)

3 Numerical Integration


We have seen that we can find the exact area under a curve by finding the definite integral. This works well as long as the function is one that we can integrate using the rules for integration or the table of integrals. This may not always be the case. The function may be one that cannot be integrated, or it may be given in the form of a table of values rather than an algebraic expression. In cases like these, we can use numerical methods to find the value of the definite integral to whatever degree of accuracy we need, using a calculator or computer.

We have already seen one numerical method for approximating areas under a curve – the midpoint method. We will look at two more methods that give us more accuracy than the midpoint method for a given number of intervals – the trapezoidal rule and Simpson's rule. Of course, the accuracy of all three methods is improved by increasing the number of intervals. When the number of intervals is very large, a computer is absolutely essential.

3.1 The Trapezoidal Rule

In the diagram shown on the right, suppose we wish to estimate the area under the curve $y = f(x)$ between $x = a$ and $x = b$.

We have divided the interval from $x = a$ to $x = b$ into n equal intervals of width h . If we join the tops of the vertical lines by straight line segments, we form trapezoids each of whose areas is *approximately* equal to the area under the curve between successive x -values.



The height of each trapezoid (measured horizontally) is equal to h , and the lengths of the parallel sides are equal to the y -values on either side of the trapezoid. Making use of the fact that the area of a trapezoid is equal to half the height times the sum of the parallel sides, we get the following expression for the sum of the areas of the trapezoids between a and b :

$$A_T = \frac{1}{2}h(y_0 + y_1) + \frac{1}{2}h(y_1 + y_2) + \frac{1}{2}h(y_2 + y_3) + \cdots + \frac{1}{2}h(y_{n-2} + y_{n-1}) + \frac{1}{2}h(y_{n-1} + y_n)$$

Factoring out the common factor $\frac{1}{2}h$ and collecting like terms, we get

$$\begin{aligned} A_T &= \frac{h}{2} [y_0 + y_1 + y_1 + y_2 + y_2 + \cdots + y_{n-1} + y_{n-1} + y_n] \\ &= \frac{h}{2} [y_0 + 2y_1 + 2y_2 + \cdots + 2y_{n-1} + y_n] \end{aligned}$$

Since the sum of the areas of the trapezoids is approximately equal to the area under the curve from a to b , which is in turn equal to the definite integral from a to b , we can state the following formula, called the *trapezoidal rule*:

$$\int_a^b f(x) dx \approx \frac{h}{2} [y_0 + 2(y_1 + y_2 + \cdots + y_{n-1}) + y_n] \quad (4)$$

where $h = \frac{b-a}{n}$ and n is the number of intervals used.

It should be fairly obvious that, as the number of intervals is increased and the width of the intervals is decreased, the area of each trapezoid approaches the area under the curve for each interval more and more closely. Thus any desired degree of accuracy can be obtained simply by using a sufficiently large number of intervals.

It should be noted that the trapezoidal rule can be used whether the function is given as a mathematical equation or as a table of values. Also, it can be used to evaluate a definite integral even for a function that cannot be integrated analytically.

Examples

- Evaluate the definite integral $\int_0^{10} x^{\frac{3}{2}} dx$
 1. using the trapezoidal rule
 - (a) with 5 panels of equal width
 - (b) with 10 panels of equal width
 2. by integration

Carry 6 significant digits and round your answers to 4 significant digits. Compare each of your results for part (a) with the answer found in part (b).

Additional Examples (Text, p. 1000)

- Problem 2
- Problem 6
- Problem 10
- Problem 14

Reading:

Sec. 34-9, pp. 996 to 998

Problems:

Ex. 9, p. 1000 # 1-17 (odd) – use trapezoidal rule only

3.2 Simpson's Rule

In the previous section, we approximated the area under a curve by summing the areas of trapezoids formed by connecting successive points of the curve with straight-line segments. Suppose that, instead of straight-line segments, we connect each successive set of three points on the curve with a parabolic arc. This leads to a formula called *Simpson's Rule*, which is derived in the text (pp. 998-1000). According to Simpson's Rule, the definite integral of a function between two points is given by the following formula:

$$\int_a^b f(x) dx \approx \frac{h}{3} [y_0 + 4y_1 + 2y_2 + \cdots + 2y_{n-2} + 4y_{n-1} + y_n] \quad (5)$$

where $h = \frac{b-a}{n}$ is the width of each panel and n is the number of panels.

It is intuitively clear that, by using parabolic arcs instead of straight line segments to join the points on a curve, we should get a better approximation of the area under a curve for the same number of intervals.

It should be noted that, in order to apply Simpson's Rule, we must have an *even* number of intervals.

As was the case with the trapezoidal rule, using more intervals leads to higher accuracy.

Example

- Evaluate the definite integral $\int_0^{10} x^{\frac{3}{2}} dx$ using Simpson's rule with 10 intervals. Carry 6 significant digits and round your answer to 4 significant digits. Compare your answer to that found on the previous page using the trapezoidal rule, and to the answer found by integration.

Additional Examples (Text, p. 1000)

- Problem 12
- Problem 16

Reading:

Sec 34-9, pp. 998-1000

Problems:

Ex. 9, p. 1000 # 1-17 (odd) – use Simpson's rule only

4 Review Problems: The Definite Integral

Questions:

1. Evaluate the following definite integrals. Give the answers in exact form and in approximate decimal form to three significant digits.

(a) $\int_1^3 x^2(3x^3 - 1)^2 dx$

(b) $\int_0^1 \frac{t dt}{4 + t^2}$

2. Given the function $y = 2x^3 + 5$,

(a) Use *Simpson's rule* with 4 intervals to find the approximate area under the curve between $x = 0$ and $x = 4$.

(b) Find the *exact* area under the above curve from $x = 0$ to $x = 4$ by integration.

(Note: If your answers to (a) and (b) are not reasonably close to each other, you'd better recheck your work!)

3. Use the *trapezoidal rule* to find the approximate area under the curve defined by the following set of experimental data:

x	5.0	6.0	7.0	8.0	9.0	10.0
y	4.2	3.9	3.8	4.0	3.5	3.4

4. Additional Review Problems:

- Chapter 30 Review Problems (p. 883) # 10-14 (all) , 28, 29
- Chapter 34 Review Problems (p. 1002) # 21 (use the trapezoidal rule with 8 intervals)

Answers:

1. (a) $\frac{56,888}{3} \approx 1.90 \times 10^4$
(b) $\frac{1}{2}(\ln 5 - \ln 4) \approx 0.112$
2. (a) 148 square units
(b) 148 square units
3. 19 square units
4. (Answers to Even-Numbered Review Problems):
 - Chapter 30:
 10. $\ln 2 \approx 0.693$
 12. $\frac{\ln 2}{2} \approx 0.347$
 14. $-\frac{14}{9} \approx -1.56$
 28. 280

5 Formulas

The Definite Integral

$$\int_a^b f(x) \, dx = F(b) - F(a) \quad (1)$$

where F is the antiderivative (indefinite integral) of f .

$$A_{ab} = \int_a^b f(x) \, dx \quad (2)$$

where A_{ab} is the area under the curve $f(x)$ between $x = a$ and $x = b$.

Numerical Approximations of the Definite Integral

Midpoint Method:

$$\int_a^b f(x) \, dx \approx \sum_{i=1}^n f(x_i^*) \Delta x = f(x_1^*) \Delta x + \cdots + f(x_n^*) \Delta x \quad (3)$$

where $f(x_i^*)$ is the value of the function at the midpoint x_i^* of each panel, n is the number of panels, and Δx is the width of each panel, $\Delta x = \frac{b-a}{n}$.

Trapezoidal rule:

$$\int_a^b f(x) \, dx \approx \frac{h}{2} [y_0 + 2(y_1 + y_2 + \cdots + y_{n-1}) + y_n] \quad (4)$$

where $h = \frac{b-a}{n}$, n is the number of intervals used, and $y_i = f(x_i)$.

Simpson's Rule:

$$\int_a^b f(x) \, dx \approx \frac{h}{3} [y_0 + 4y_1 + 2y_2 + \cdots + 2y_{n-2} + 4y_{n-1} + y_n] \quad (5)$$

where $h = \frac{b-a}{n}$ is the width of each panel, n is the (even) number of panels, and $y_i = f(x_i)$.

Version 1.3, 3 November 2008
Copyright © 2000, 2001, 2002, 2007, 2008 Free Software Foundation, Inc.
<<http://fsf.org/>>
Everyone is permitted to copy and distribute verbatim copies of this license document, but changing it is not allowed.

You may copy and distribute the Document in any medium, either commercially or noncommercially, provided that this License, the copyright notices, and the license notice saying this License applies to the Document are reproduced in all copies. You may not use technical measures to obstruct or control the reading or use of this License. You may not use technical measures to obstruct or control the reading or use of copies you make or distribute. However, you may accept compensation in exchange for copies. If you distribute a large enough number of copies you must also follow the conditions in section 3.

You may charge a fee for copying, distributing, or performing the Document under this License, provided you state on the copies, under these conditions stated above, and you prominently display copies of the License and the copyright notices, and you

3. COPYING IN QUANTITY

[illegible]

This License applies to any manual or other work, in any medium, that contains a notice placed by the copyright holder saying it can be distributed under the terms of this License. Such a notice grants a world-wide, royalty-free license, and no additional restrictions may be placed on the exercise of these rights. For example, "Document" below refers to any such manual or work. Any number of the public is a licensee, and is addressed as "you". You accept the license if you copy, modify or distribute the work in a way requiring permission under copyright law. A "Modified Version" of the Document means any work containing the Document or a portion of it, with modifications to it, made by you or by others acting under your authority to produce modified versions. Such modified versions must be clearly marked as such, and must not be presented as being the original work in any medium. You may create a modified version, provided that you release it under its own license, which must contain, in the same form and at least as free as the original license, the following notice:

[illegible]

The “**Title Page**” means, for a printed book, the title page itself, plus such following pages as are needed to hold, legibly, the material this License requires to appear in the title page. For works in formats which do not have any title page as such, “Title Page” means the text near the most prominent appearance of the work’s title, preceding the beginning of the body of the text.

The Document may include Warranty Disclaimers next to the notice which states that this License applies to the Document. These Warranty Disclaimers are considered to be included by reference in this License, but only as regards disclaiming warranties; any other implication that these Warranty Disclaimers may have is void and has no effect on the meaning of this License.

You may copy and distribute a Modified Version of the Document under the conditions of sections 2 and 3 above, provided that you release the Modified Version under precisely this License, with the Modified Version filling the role of the Document, thus licensing distribution and modification of the Modified Version to whoever possesses a copy of it. In addition, you must do these things to the Modified Version:

F. Include, immediately after the copyright notice, a license notice giving the public permission to use the Modified Version under the terms of this License, in the form shown in the Addendum below.

J. Preserve the network location, i.e., given in the Document for public access to a Transparent copy of the Document, and likewise the network locations given in the Document for previous versions if it was based on them. These may be placed in the "History" section. You may omit a network location for a work that was published at least four years before the Document version itself, or if the original publisher of the version it refers to gives permission.

M. Delete any section Entitled "Endorsements". Such a section may not be included in the Modified Version.

N. Do not retile any existing section to be Entitled "Endorsements" or to conflict in title with any Invariant Section.

O. Preserve any Warranty Disclaimers.

You may make a collection consisting of the Document and other documents released under this License, and replace the individual copies of this License in the released documents with a single copy that is included in the collection, provided that you follow the rules of this License for verbatim copying of each of the documents in all other respects.

You may extract a single document from such a collection, and distribute it individually under this License, provided you insert a copy of this License into the new document, and follow the rules of this License in all other respects regarding verbatim copying of that document.

WORKS

8. TRANSLATION

[illegible]

However, if you cause a violation of this License, then your license from a particular copyright holder is terminated (a) provisionally, unless and until the copyright holder notifies you of the alleged violation by a reasonable means, and you cease the violation; and (b) permanently, if the copyright holder fails to notify you of the violation by some reasonable means prior to 60 days after the cessation.

The Free Software Foundation may publish new, revised versions of the GNU Free Documentation License from time to time. Such new versions will be similar in spirit to the present version, but may differ in detail to address new problems or concerns. See <http://www.gnu.org/copyleft/>.

11. RELICENSING

An MMC is "eligible for relicensing" if it is licensed under this License, and if all works that were first published under this License somewhere other than this MMC, and subsequently incorporated in whole or in part into the MMC, (1) had no cover texts or invariant sections, and (2) were thus incorporated prior to November 1, 2008.

**ADDENDUM: How to use this License for
your documents**

Copyright © YEAR YOUR NAME. Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.3 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled "GNU Free Documentation License".

with the Invariant Sections being LIST THEIR TITLES, with the Front-Cover Texts being LIST, and with the Back-Cover Texts being LIST.

If you have Invariant Sections without Cover Texts, or some other combination of the three, merge those two alternatives to suit the situation.

If your document contains nontrivial examples of program code, we recommend releasing these examples in parallel under your choice of free software license, such as the GNU General Public License, to permit their use in free software.